arXiv:q-alg/9712055v2 12 Jan 1998

ITEP-TH-68/97

The q-Fourier transform of q-distributions.

M.A.Olshanetsky
ITEP, 117259, Moscow
e-mail olshanet@heron.itep.ru

V.-B.K.Rogov
MIIT, 101475, Moscow
e-mail m10106@sucemi.bitnet

Abstract

We consider functions on the lattice generated by the integer powers of q^2 for $0 < q < 1$ and construct the q-analog of Fourier transform based on the Jackson integral in the space of distributions on the lattice.

1 Introduction

The classical Fourier transform is a powerful tool in the Harmonic analysis on the simple Lie groups and homogeneous spaces. In the quantum group case similar problems can be simplified by the Fourier transform in the quantum affine space. There exist a few papers devoted to the quantum Fourier transform [1, 2], but their main object is the quantum algebra of operators without specification of the spaces of functions where these operators are defined. For this reason, applications of those results in concrete problems of Harmonic analysis are not so easy. We investigate here the simplest one-dimensional case where the whole effect of the non-commutativity emerges only in the lattice form of the space of function and a very simple algebra generated by functions in the direct and inverse Fourier spaces. This particular situation can be considered as the q-deformation of the classical Fourier transform which attracted last time a particular attention due its relation to the generalizations of the classical harmonic oscillator problem (see [3, 4, 5, 6]). While one of the primary interest of these papers is the investigations of different kernels of the q-Fourier transforms, we focus on spaces of functions and their special q-Fourier transform based on the q-exponents. Our approach is very similar to the classical picture developed in [7]. We consider the space of test functions and q^2-distributions and then determine the q^2-Fourier transform of test functions. A slightly different q-Fourier transform on a subspace of test functions was considered in [8]. The q^2-Fourier transform of the q^2-distributions is determined by the Parseval identity. In conclusion, we present a table of the q^2-Fourier transforms of some q^2-distributions.

2 Preliminary relations

We assume that $z \in \mathbf{C}$ and $|q| < 1$, unless otherwise is specified.

We recall some notations [9]. For an arbitrary a

$$(a,q)_n = \begin{cases} 1 & for \quad n = 0 \\ (1-a)(1-aq)\dots(1-aq^{n-1}) & for \quad n \geq 1, \end{cases}$$

$$(a,q)_\infty = \lim_{n\to\infty}(a,q)_n.$$

$$\begin{bmatrix} l \\ i \end{bmatrix}_{q^2} = \frac{(q^2,q^2)_l}{(q^2,q^2)_i(q^2,q^2)_{l-i}}.$$

Consider the q^2-exponentials

$$e_{q^2}(z) = \sum_{n=0}^{\infty} \frac{z^n}{(q^2,q^2)_n} = \frac{1}{(z,q^2)_\infty}, \qquad |z| < 1, \tag{2.1}$$

$$E_{q^2}(z) = \sum_{n=0}^{\infty} \frac{q^{n(n-1)}z^n}{(q^2,q^2)_n} = (-z,q^2)_\infty \tag{2.2}$$

and the q^2-trigonometric functions

$$\cos_{q^2} z = \frac{1}{2}(e_{q^2}(iz) + e_{q^2}(-iz)), \quad \sin_{q^2} z = \frac{1}{2i}(e_{q^2}(iz) - e_{q^2}(-iz)), \tag{2.3}$$

$$\mathrm{Cos}_{q^2} z = \frac{1}{2}(E_{q^2}(iz) + E_{q^2}(-iz)), \quad \mathrm{Sin}_{q^2} z = \frac{1}{2i}(E_{q^2}(iz) - E_{q^2}(-iz)). \tag{2.4}$$

We also consider the basic hypergeometric series

$${}_0\Phi_1(-;0;q^2,z) = \sum_{n=0}^{\infty} \frac{q^{2n(n-1)}z^n}{(q^2,q^2)_n}. \tag{2.5}$$

It follows from (2.1) that $e_{q^2}(z)$ is a meromorphic function with simple poles in the points $z = q^{-2k}, k = 0, 1, \dots$.

Lemma 2.1 *The function $e_{q^2}(z)$ is represented as the sum of partial fractions:*

$$e_{q^2}(z) = \frac{1}{(q^2,q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k(1-zq^{2k})}. \tag{2.6}$$

Proof. Let

$$e_{q^2}(z,n) = \frac{1}{(z,q^2)_n} = \sum_{k=0}^{n} \frac{c_{k,n}}{1-zq^{2k}},$$

where

$$c_{k,n} = \mathrm{res}_{z=q^{-2k}}\, e_{q^2}(z,n) = \lim_{z\to q^{-2k}} (1-zq^{2k})e_{q^2}(z,n) =$$

$$= \frac{1}{(1-q^{-2k})\dots(1-q^{-2})(1-q^2)\dots(1-q^{2(n-k)})} = \frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k (q^2,q^2)_{n-k}}.$$

Therefore,

$$e_{q^2}(z,n) = \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k (q^2,q^2)_{n-k}(1-aq^{2k})}.$$

From $\frac{1}{(q^2,q^2)_{n-k}} < \frac{1}{(q^2,q^2)_\infty} = e_{q^2}(q^2)$, we obtain

$$e_{q^2}(z) = \lim_{n\to\infty} e_{q^2}(z,n) = \frac{1}{(q^2,q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k (1-zq^{2k})}.$$

Since this series converges absolutely for $q \in (0,1)$, (2.6) holds for an arbitrary $z \neq q^{-2k}$. ■

From (2.3) we obtain

Corollary 2.1 *The functions* $\cos_{q^2} z$ *and* $\sin_{q^2} z$ *are represented as the sums of partial fractions*

$$\cos_{q^2} z = \frac{1}{(q^2,q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k (1+z^2q^{4k})}, \tag{2.7}$$

$$\sin_{q^2} z = \frac{z}{(q^2,q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+3)}}{(q^2,q^2)_k (1+z^2q^{4k})}. \tag{2.8}$$

Proposition 2.1 *If* z *and* q *are real then*

$$|\cos_{q^2} z| \leq \frac{(-q^2,q^2)_\infty}{(1+z^2)(q^2,q^2)_\infty},$$

$$|\sin_{q^2} z| \leq \frac{|z|(-1,q^2)_\infty}{(1+z^2)(q^2,q^2)_\infty}.$$

Proof. Since

$$\frac{1}{1+z^2q^{4k}} \leq \frac{q^{-4k}}{1+z^2},$$

for an arbitrary real z and for an arbitrary $k \geq 0$, we have

$$\left|\sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k (1+z^2q^{4k})}\right| \leq \frac{1}{1+z^2} \sum_{k=0}^{\infty} \frac{q^{k(k-1)}q^{-2k}}{(q^2,q^2)_k} = \frac{1}{1+z^2} E_{q^2}(q^{-2}),$$

$$\left|\sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+3)}}{(q^2,q^2)_k (1+z^2q^{4k})}\right| \leq \frac{1}{1+z^2} \sum_{k=0}^{\infty} \frac{q^{k(k-1)}}{(q^2,q^2)_k} = \frac{1}{1+z^2} E_{q^2}(1).$$

The statement of Proposition follows from (2.7), (2.8) and the last relations. ■

The next proposition follows immediately from (2.2) and (2.4).

Proposition 2.2 $\mathrm{Cos}_{q^2} z$ *and* $\mathrm{Sin}_{q^2} z$ *satisfy the inequalities*

$$|\mathrm{Cos}_{q^2} z| \leq 1, \qquad |\mathrm{Sin}_{q^2} z| \leq |z|. \tag{2.9}$$

Consider the theta function

$$\vartheta(u|\tau) = 2p^{1/8}\sin\pi u\prod_{n>0}(1-e^{2\pi iu}p^n)(1-e^{-2\pi iu}p^n)(1-p^n)$$

where $p = \exp 2\pi i\tau$, $(Im\tau > 0)$. Let

$$\mathbf{Q}(z,q) = (1-q^2)\sum_{m=-\infty}^{\infty}\frac{1}{zq^{2m}+z^{-1}q^{-2m}} \tag{2.10}$$

Then it follows from the modular properties of (2.10) that

$$\mathbf{Q}(z,q) = -\frac{1}{2\pi i}\frac{\vartheta(u+\tau/2|\tau)\vartheta'(0|\tau)}{\vartheta(u|\tau)\vartheta(\tau/2|\tau)},$$

where

$$\tau = -\frac{2i\ln q}{\pi}, \quad u = \frac{1}{\pi i}\ln z + \frac{1}{2}.$$

Define the function

$$\Theta(z) = (1-q^2)\sum_{m=-\infty}^{\infty}\sin_{q^2}((1-q^2)q^{2m}z). \tag{2.11}$$

Proposition 2.3 *$\Theta(z)$ has the following properties:*
1. $\Theta(q^{2k}z) = \Theta(z)$ for any $z \neq 0$,
2. $\Theta(z) = \mathbf{Q}((1-q^2)z,q)$.

Proof. 1. follows immediately from (2.11).

Due to (2.8) $\Theta(z)$ can be represented as

$$\Theta(z) = \frac{1-q^2}{(q^2,q^2)_\infty}\sum_{m=-\infty}^{\infty}q^{2m}\sum_{k=0}^{\infty}\frac{(-1)^k q^{k(k+3)}(1-q^2)z}{(q^2,q^2)_k[1+(1-q^2)^2z^2q^{4(m+k)}]}.$$

We can change the order of summation since the inner series converges uniformly with respect to m. Then using (2.1), (2.2) and (2.10) we obtain

$$\Theta(z) = \frac{1-q^2}{(q^2,q^2)_\infty}\sum_{k=0}^{\infty}\frac{(-1)^k q^{k(k+3)}}{(q^2,q^2)_k}\sum_{m=-\infty}^{\infty}\frac{(1-q^2)zq^{2m}}{1+(1-q^2)^2z^2q^{4(m+k)}} =$$

$$= \frac{1-q^2}{(q^2,q^2)_\infty}\sum_{k=0}^{\infty}\frac{(-1)^k q^{k(k+1)}}{(q^2,q^2)_k}\sum_{m=-\infty}^{\infty}\frac{(1-q^2)zq^{2m}}{1+(1-q^2)^2z^2q^{4m}} = \mathbf{Q}((1-q^2)z,q).$$

■

Specially we define

$$\Theta_0 = \Theta(1) = \mathbf{Q}(1-q^2,q). \tag{2.12}$$

Proposition 2.4 *For an arbitrary integer $M > 0$*

$$(1-q^2)z\sum_{m=-M}^{\infty} q^{2m}\cos_{q^2}((1-q^2)q^{2m}z) = \sin_{q^2}((1-q^2)q^{-2M}z), \tag{2.13}$$

$$(1-q^2)z\sum_{m=-M}^{\infty} q^{2m}\sin_{q^2}((1-q^2)q^{2m}z) = 1-\cos_{q^2}((1-q^2)q^{-2M}z). \tag{2.14}$$

Proof. Assume that $|(1-q^2)q^{-2m}z| < 1$. Then

$$(1-q^2)z\sum_{m=-M}^{\infty} q^{2m}\cos_{q^2}((1-q^2)q^{2m}z) = \sum_{m=-M}^{\infty}(1-q^2)zq^{2m}\sum_{k=0}^{\infty}(-1)^k\frac{(1-q^2)^{2k}q^{4mk}z^{2k}}{(q^2,q^2)_{2k}}$$

$$= \sum_{k=0}^{\infty}(-1)^k\frac{(1-q^2)^{2k+1}z^{2k+1}}{(q^2,q^2)_{2k}}\sum_{m=-M}^{\infty}q^{2m(2k+1)} = \sin_{q^2}((1-q^2)q^{-2M}z). \tag{2.15}$$

For $z \neq \pm i(1-q^2)^{-1}q^{-2n}$ the left side of (2.13) is defined as the analytic continuation of (2.15). (2.14) can be proved in the same way. ■

3 q^2-derivatives and q^2-integral

Let $\mathcal{A} = C(z, z^{-1})$ be the algebra of formal Laurent series. The q^2-derivative of $f(z) \in \mathcal{A}$ is defined as

$$\partial_z f(z) = \frac{z^{-1}}{1-q^2}(f(z) - f(q^2z)). \tag{3.1}$$

For an arbitrary $n \geq 0$

$$\partial_z^k z^n = \begin{cases} \frac{(q^2,q^2)_n}{(q^2,q^2)_{n-k}(1-q^2)^k} z^{n-k} & for \quad 0 \leq k \leq n \\ 0 & for \quad k > n, \end{cases} \tag{3.2}$$

and for any $n \geq 0$ and $k \geq 0$

$$\partial_z^k z^{-n-1} = (-1)^k q^{-k(2n+k+1)}\frac{(q^2,q^2)_{n+k}}{(q^2,q^2)_n(1-q^2)^k}z^{-n-k-1}. \tag{3.3}$$

The q^2-integral (Jackson integral [9]) is defined as the map I_{q^2} from $\mathcal{A}$ to the space of formal number series

$$I_{q^2}f = \int d_{q^2}zf(z) = (1-q^2)\sum_{m=-\infty}^{\infty} q^{2m}[f(q^{2m}) + f(-q^{2m})] \tag{3.4}$$

Definition 3.1 *The function $f(z)$ is locally q^2-integrable, if the q^2-integral*

$$\int_a^b d_{q^2}zf(z) = (1-q^2)\sum_{m=0}^{\infty} q^{2m}[bf(bq^{2m}) - af(aq^{2m})] \tag{3.5}$$

exists for any finite a and b, i.e., the series in the right side of (3.5) converges.

Definition 3.2 *$f(z)$ is absolutely q^2-integrable function, if the series*

$$\sum_{m=-\infty}^{\infty} q^{2m}[|f(q^{2m})| + |f(-q^{2m})|]$$

converges.

Let $\mathcal{B}$ be the algebra analogous to $\mathcal{A}$, but generated by s, s^{-1} which commute with z as $zs = q^2 sz$. Define the q^2-differentiation in $\mathcal{B}$ as

$$\partial_s \phi(s) = (\phi(s) - \phi(q^2 s))\frac{s^{-1}}{1-q^2}.$$

We denote by $\mathcal{AB}$ the whole algebra with two generators z, z^{-1}, s, s^{-1} and the q^2-differentiation

$$zs = q^2 sz, \quad \partial_z s = q^{-2} s\partial_z, \quad \partial_s z = q^2 z\partial_s, \quad \partial_z\partial_s = q^2\partial_s\partial_z. \tag{3.6}$$

We consider $\mathcal{AB}$ as a left module under the left action of $\mathcal{A}$ by multiplication, and a right module under the left action of $\mathcal{B}$.

To define the q^2-integral on $\mathcal{AB}$ we order the generators of integrand in such a way that z stays on the left side while s stays on the right side. For example, if $f(z) = \sum_r a_r z^r$ then

$$f(zs) = \sum_r a_r (zs)^r = \sum_r a_r q^{-r(r-1)} z^r s^r.$$

For convenience we introduce the following notation

$$\ddagger g(zs)\ddagger = \sum_r a_r z^r s^r, \text{ if } g(z) = \sum_r a_r z^r.$$

For example, we can derive from (2.1), (2.2) and (3.6) that

$$E_{q^2}(i(1-q^2)zs) = \ddagger e_{q^2}(i(1-q^2)zs)\ddagger.$$

Following prescribed rules we calculate
1.

$$\int d_{q^2} z z^{-1} E_{q^2}(i(1-q^2)zs) = 2i\Theta(s) \tag{3.7}$$

(see (2.12)).
2. Using (2.14) and Proposition 2.1 we get

$$\int d_{q^2} z \Theta(z) E_{q^2}(i(1-q^2)zs) = 2i\Theta_0 s^{-1}.$$

4 q^2-distributions

Let $S_{q^2} = \{\phi(x)\}$ be the space of infinitely q^2-differentiable fast decreasing functions

$$|x^k \partial_x^l \phi(x)| \le C_{k,l}(q), \ k \ge 0, \ l \ge 0. \tag{4.1}$$

Let S be the space of infinitely differentiable (in the classical sense) fast decreasing functions

$$|x^k \phi^{(l)}(x)| \le C_{k,l}, \ k \ge 0 \ l \ge 0. \tag{4.2}$$

Proposition 4.1

$$S \subset S_{q^2}.$$

Proof. We denote by $\phi(x_1, x_2, \ldots, x_k)$ the separated difference of order k. Then it is easy to find that for an arbitrary integer $l \geq 0$

$$\phi(q^{2l}x, q^{2l-2}x, \ldots, x) = \frac{(q^2, q^2)_l}{(1-q^2)^l} \partial_x^l \phi(x). \tag{4.3}$$

On the other hand, if $\phi(x)$ is differentiable l times (in the classical sense), then there exists a point $\xi \in (q^{2l}x, x)$, such that

$$\phi(q^{2l}x, q^{2l-2}x, \ldots, x) = \frac{1}{l!} \phi^{(l)}(\xi). \tag{4.4}$$

Then, it follows from (4.3) and (4.4)

$$\partial_x^l \phi(x) = \frac{(1-q^2)^l}{(q^2, q^2)_l l!} \phi^{(l)}(\xi), \quad \xi \in (q^{2l}x, x).$$

Therefore, from (4.2)

$$|x^k \partial_x^l \phi(x)| \leq \frac{(1-q^2)^l}{(q^2, q^2)_l l!} C_{k,l}.$$

Finally, if $\phi_n(x) \to 0$ for $n \to \infty$ in the space S then $\phi_n(x) \to 0$ in the space S_{q^2}. ■

Definition 4.1 *The skeleton $\hat{\phi}(z)$ of $\phi(z) \in S_{q^2}$ is the set of evaluations of $\phi(z)$ on the lattice generated by the powers of q^2*

$$\hat{\phi}(z) = \phi(z)|_{z=q^{2n}}, n = 0, \pm 1, \ldots.$$

The space of the skeletons we denote by $\hat{S}_{q^2}$.

The functions

$$\hat{\phi}_n^+(z) = \begin{cases} 1 & \text{for} \quad z = q^{2n} \\ 0 & \text{for} \quad z \neq q^{2n}, \end{cases} \quad \hat{\phi}_n^-(z) = \begin{cases} 1 & \text{for} \quad z = -q^{2n} \\ 0 & \text{for} \quad z \neq -q^{2n}. \end{cases}$$

generate the basis in the space $\hat{S}_{q^2}$:

$$\hat{\phi}(z) = \sum_{n=-\infty}^{\infty} [a_n \hat{\phi}_n^+(z) + b_n \hat{\phi}_n^-(z)], \quad a_n = \phi(q^{2n}), \quad b_n = \phi(-q^{2n}).$$

The topology in $\hat{S}_{q^2}$ is induced by the topology in S_{q^2} (4.1):

$$|\frac{q^{2n(k-l)}}{(1-q^2)^l} \sum_{i=1}^{l} (-1)^i \left[\begin{array}{c} l \\ i \end{array} \right]_{q^2} q^{-i(2l-i-1)} a_{n+i}| \leq C_{k,l}(q),$$

$$|(-1)^{k-l} \frac{q^{2n(k-l)}}{(1-q^2)^l} \sum_{i=1}^{l} (-1)^i \left[\begin{array}{c} l \\ i \end{array} \right]_{q^2} q^{-i(2l-i-1)} b_{n+i}| \leq C_{k,l}(q).$$

Let L_{q^2} be the linear map from S_{q^2} to $\hat{S}_{q^2}$ defined by the evaluation of functions in the vertices of the lattice. Then

$$\hat{S}_{q^2} = S_{q^2} / \ker L_{q^2}.$$

Let $\Lambda\phi(z) = \phi(q^2 z)$. Then

$$\Lambda z = q^2 z \Lambda, \qquad \partial_z \Lambda = q^2 \Lambda \partial_z. \tag{4.5}$$

The operations Λ and ∂_z are well defined on $\hat{S}_{q^2}$ since

$$\Lambda L_{q^2} = L_{q^2} \Lambda, \quad \partial_z L_{q^2} = L_{q^2} \partial_z.$$

Moreover, the q^2-integral vanishes on the functions from $\ker L_{q^2}$, and therefore is defined on the quotient $\hat{S}_{q^2} = S_{q^2} / \ker L_{q^2}$. In some places, we will write in the q^2-integral an element from $\hat{S}_{q^2}$ assuming that it is a representative from the quotient.

Proposition 4.2 *If* $\hat{\phi}(z) \in \hat{S}_{q^2}$ *then*

$$\int d_{q^2} z \partial_z \hat{\phi}(z) = 0.$$

Proof. It follows from (3.1),(3.4) and (4.1)

$$\int d_{q^2} z \partial_z \hat{\phi}(z) = \sum_{m=-\infty}^{\infty} [\hat{\phi}(q^{2m}) - \hat{\phi}(q^{2m+2}) - \hat{\phi}(-q^{2m}) + \hat{\phi}(-q^{2m+2})] =$$

$$= \lim_{M \to \infty} [\hat{\phi}(q^{-2M}) - \hat{\phi}(-q^{-2M})] = 0.$$

■

Corollary 4.1 *(q^2-integration by parts) For any* $k \geq 0$

$$\int d_{q^2} z \hat{\phi}(z) \partial_z^k \hat{\psi}(z) = (-1)^k q^{-k(k-1)} \int d_{q^2} z \partial_z^k \hat{\phi}(z) \hat{\psi}(q^{2k} z).$$

Definition 4.2 *The q^2-distribution f over $\hat{S}_{q^2}$ is a linear continuous functional* $f : \hat{S}_{q^2} \to \mathbf{C}$.

We denote by $\hat{S}'_{q^2}$ the space of the q^2-distributions over $\hat{S}_{q^2}$.

Definition 4.3 *The sequence $\{f_n\}$ is referred to as convergent in $\hat{S}'_{q^2}$ to $f \in \hat{S}'_{q^2}$, if for any $\phi(z) \in \hat{S}_{q^2}$ the sequence $\{< f_n, \phi >\}$ converges to $< f, \phi >$.*

The q^2-distributions which are defined by the q^2-integral

$$< f, \phi >= \int_{\infty}^{\infty} d_{q^2} z \bar{f}(z) \phi(z) = (1 - q^2) \sum_{m=-\infty}^{\infty} q^{2m} [\bar{f}(q^{2m}) \phi(q^{2m}) + \bar{f}(-q^{2m}) \phi(-q^{2m})]$$

we refer as regular.

Corollary 4.1 and Proposition 4.2 allows to introduce the q^2-differentiation in $\hat{S}'_{q^2}$

$$< \partial_z f, \phi >= -q^2 < \Lambda f, \partial_z \phi > . \tag{4.6}$$

Let us give some examples of q^2-distributions.

1.
$$< \theta^+_{q^2}, \phi >= \int_0^\infty d_{q^2} z \hat{\phi}(z) = (1-q^2) \sum_{m=-\infty}^{\infty} q^{2m} \phi(q^{2m}). \tag{4.7}$$

2.
$$< \theta^-_{q^2}, \phi >= \int_{-\infty}^0 d_{q^2} z \hat{\phi}(z) = (1-q^2) \sum_{m=-\infty}^{\infty} q^{2m} \phi(-q^{2m}). \tag{4.8}$$

So, the q^2-distributions $\theta^+_{q^2}$ and $\theta^-_{q^2}$ correspond to the functions

$$\hat{\theta}^+(z) = \sum_{n=-\infty}^{\infty} \hat{\phi}^+_n(z), \qquad \hat{\theta}^-(z) = \sum_{n=-\infty}^{\infty} \hat{\phi}^-_n(z).$$

3.
$$< \delta_{q^2}, \phi >= \phi(0) = \lim_{m \to \infty} \frac{\phi(q^{2m}) + \phi(-q^{2m})}{2}.$$

Moreover, it follows from (4.6)

$$< \partial_z (\theta^+_{q^2} - \theta^-_{q^2}), \phi >= - < \theta^+_{q^2} - \theta^-_{q^2}, \partial_z \phi >=$$

$$= -(1-q^2) \sum_{m=-\infty}^{\infty} q^{2m} \Big(\frac{q^{-2m}}{1-q^2} [\phi(q^{2m}) - \phi(q^{2m+2})] + \frac{q^{-2m}}{1-q^2} [\phi(-q^{2m}) - \phi(-q^{2m+2})] \Big) =$$

$$= - \lim_{M \to \infty} \sum_{m=-M}^{M} [\phi(q^{2m}) - \phi(q^{2m+2}) + \phi(-q^{2m}) - \phi(-q^{2m+2})] =$$

$$= - \lim_{M \to \infty} [\phi(q^{-2M}) - \phi(q^{2M+2}) + \phi(-q^{-2M}) - \phi(-q^{2M+2})] = 2\phi(0),$$

i.e.,

$$\partial_z (\theta^+_{q^2}(z) - \theta^-_{q^2}(z)) = 2\delta_{q^2}(z). \tag{4.9}$$

4.
$$< z^{-1}, \phi >= \int_0^\infty d_{q^2} z z^{-1} [\hat{\phi}(z) - \hat{\phi}(-z)] = (1-q^2) \sum_{m=-\infty}^{\infty} [\phi(q^{2m}) - \phi(-q^{2m})].$$

5. For an arbitrary $k \geq 0$

$$< z^{-k-1}, \phi >= (-1)^k q^{k(k+1)} \frac{(1-q^2)^k}{(q^2,q^2)_k} < \partial_z^k z^{-1}, \phi >= \frac{(1-q^2)^k}{(q^2,q^2)_k} < z^{-1}, \partial_z^k \phi >=$$

$$= \frac{(1-q^2)^{k+1}}{(q^2,q^2)_k} \sum_{m=-\infty}^{\infty} [\partial_z^k \phi(z)|_{z=q^{2m}} - \partial_z^k \phi(z)|_{z=-q^{2m}}]. \tag{4.10}$$

6. For an arbitrary $\nu > -1$

$$< z_+^\nu, \phi >= \int_0^\infty d_{q^2} z z^\nu \hat{\phi}(z) = (1-q^2) \sum_{m=-\infty}^{\infty} q^{2m(\nu+1)} \phi(q^{2m}).$$

Since for any $k \geq 0$

$$\partial_z^k z^\nu = (-1)^k q^{k(2\nu-k+1)} \frac{(q^{-2\nu}, q^2)_k}{(1-q^2)^k} z^{\nu-k},$$

we define

$$< z_+^{\nu-k}, \phi >= \frac{(1-q^2)^{k+1}}{(q^{-2\nu}, q^2)_k} \sum_{m=-\infty}^{\infty} q^{2m(\nu+1)} \partial_z^k \phi(z)|_{z=q^{2m}}.$$

7. Similarly, for an arbitrary $\nu > -1$

$$< z_-^\nu, \phi >= \int_{-\infty}^0 d_{q^2} z (-z)^\nu \hat{\phi}(z) = (1-q^2) \sum_{m=-\infty}^{\infty} q^{2m(\nu+1)} \phi(-q^{2m}),$$

and for an arbitrary $k \geq 0$

$$< z_-^{\nu-k}, \phi >= (-1)^k \frac{(1-q^2)^{k+1}}{(q^{-2\nu}, q^2)_k} \sum_{m=-\infty}^{\infty} q^{2m(\nu+1)} \partial_z^k \phi(z)|_{z=-q^{2m}}.$$

5 The q^2-Fourier transform

Consider the space $S^{q^2} = \{\psi(s)\}$ generated by the similar class of functions as $S_{q^2} = \{\phi(z)\}$ (4.1), but s and z are related as the generators in the algebra $\mathcal{AB}$ (3.6). We introduce the same topology in S^{q^2} as in S_{q^2}

$$|s^k \partial_s^l \phi(s)| \leq C_{k,l}(q),\ k \geq 0,\ l \geq 0, \tag{5.1}$$

and, thereby, these spaces are isomorphic.

We define the map from S_{q^2} to S^{q^2}

$$S_{q^2} \ \xrightarrow{L_{q^2}}\ \hat{S}_{q^2} \ \xrightarrow{\mathcal{F}_{q^2}}\ S^{q^2},$$

where

$$\mathcal{F}_{q^2} \phi(z) = \int d_{q^2} z \phi(z)\ {}_0\Phi_1(-;0;q^2, i(1-q^2)q^2 zs), \tag{5.2}$$

is the q^2-Fourier transform and ${}_0\Phi_1$ is defined by (2.5). It follows from (5.2) that $\mathcal{F}_{q^2}$ is well-defined on S_{q^2}, and, in some places, we will preserve the same notation for the operator acting from S_{q^2} to S^{q^2} discarding the action of L_{q^2}.

Our aim is to find the inverse map and to prove the continuity of the both maps. More precisely, we define the dual space of skeletons $\hat{S}^{q^2}$ for S^{q^2} and the map

$$\mathcal{F}_{q^2}^{-1} \psi(s) = \frac{1}{2\Theta_0} \int E_{q^2}(-i(1-q^2)zs) \psi(s) d_{q^2} s, \quad \psi(s) \in \hat{S}^{q^2} \tag{5.3}$$

from $\hat{S}^{q^2}$ to S_{q^2}.

Consider the diagram

$$\begin{tikzcd}
S_{q^2} \arrow[r, "L_{q^2}"] & \hat{S}_{q^2} \arrow[d, "\mathcal{F}_{q^2}"] \\
\hat{S}^{q^2} \arrow[u, "\mathcal{F}^{-1}_{q^2}"] & S^{q^2}. \arrow[l, "L_{q^2}"']
\end{tikzcd}$$

We will prove

Proposition 5.1 *1.The maps*

$$\mathcal{F}_{q^2}L_{q^2} : S_{q^2} \rightarrow S^{q^2}$$

$$\mathcal{F}^{-1}_{q^2}L_{q^2} : S^{q^2} \rightarrow S_{q^2}$$

realize isomorphisms of topological spaces.
2.The maps

$$L_{q^2}\mathcal{F}^{-1}_{q^2}L_{q^2}\mathcal{F}_{q^2} : \hat{S}_{q^2} \rightarrow \hat{S}_{q^2},$$

$$L_{q^2}\mathcal{F}_{q^2}L_{q^2}\mathcal{F}^{-1}_{q^2}$$

are the identity maps on $\hat{S}_{q^2}$ *and* $\hat{S}^{q^2}$ *correspondingly.*

We begin with

Lemma 5.1

$$\int d_{q^2}z e_{q^2}(-i(1-q^2)z)\ {}_0\Phi_1(-;0;q^2,i(1-q^2)q^2zs) = \begin{cases} \frac{2}{1-q^2}\Theta_0 & for \quad s=1 \\ 0 & for \quad s\neq 1. \end{cases}$$

where Θ_0 *is determined by (2.12).*

Proof. In according with our definition of q^2-integrals all integrands must by ordered. It follows from (2.2), (2.5) and (3.6) that

$${}_0\Phi_1(-;0;q^2,i(1-q^2)q^2zs) = \ddagger E_{q^2}(i(1-q^2)q^2zs)\ddagger,$$

and we come to the integral

$$\int d_{q^2}z e_{q^2}(-i(1-q^2)z)\ddagger E_{q^2}(i(1-q^2)q^2zs)\ddagger.$$

Using (2.1) and (2.2) we obtain

$$\partial_z[e_{q^2}(-i(1-q^2)z)\ddagger E_{q^2}(i(1-q^2)zs)\ddagger] = -ie_{q^2}(-i(1-q^2)z)\ddagger E_{q^2}(i(1-q^2)q^2zs)\ddagger(1-s).$$

Hence, if $s\neq 1$, then

$$\int d_{q^2}z e_{q^2}(-i(1-q^2)z)\ddagger E_{q^2}(i(1-q^2)q^2zs)\ddagger =$$

$$= i(1-s)^{-1}\int d_{q^2}z\partial_z[e_{q^2}(-i(1-q^2)z)\ddagger E_{q^2}(i(1-q^2)zs)\ddagger].$$

It can be represented in the form (see(2.3) and (2.4))

$$(s-1)^{-1}\lim_{M\to\infty}[\cos_{q^2}((1-q^2)q^{-2M})\,\mathrm{Sin}_{q^2}((1-q^2)q^{-2M+2}s)-$$

$$-\sin_{q^2}((1-q^2)q^{-2M})\,\mathrm{Cos}_{q^2}((1-q^2)q^{-2M+2}s)-\cos_{q^2}((1-q^2)q^{2M+2})\,\mathrm{Sin}_{q^2}((1-q^2)q^{2M+4}s)+$$

$$+\sin_{q^2}((1-q^2)q^{2M+2})\,\mathrm{Cos}_{q^2}((1-q^2)q^{2M+4}s)].$$

Due to Propositions 2.1 and 2.2 the last expression vanishes.

If $s=1$ then

$$\int d_{q^2}z e_{q^2}(-i(1-q^2)z)E_{q^2}(i(1-q^2)q^2z)=$$

$$=(1-q^2)\sum_{m=-\infty}^{\infty}q^{2m}\left[\frac{(-i(1-q^2)q^{2m+2},q^2)_\infty}{(-i(1-q^2)q^{2m},q^2)_\infty}+\frac{(i(1-q^2)q^{2m+2},q^2)_\infty}{(i(1-q^2)q^{2m},q^2)_\infty}\right]=$$

$$=(1-q^2)\sum_{m=-\infty}^{\infty}q^{2m}\left(\frac{1}{1+i(1-q^2)q^{2m}}+\frac{1}{1-i(1-q^2)q^{2m}}\right)=$$

$$=2\sum_{m=\infty}^{\infty}\frac{1}{(1-q^2)^{-1}q^{-2m}+(1-q^2)q^{2m}}=\frac{2}{1-q^2}\Theta_0$$

(see (2.10) and (2.12)).■

The next Lemma can be proved in the same way.

Lemma 5.2

$$\int E_{q^2}(-i(1-q^2)zs)E_{q^2}(i(1-q^2)q^2s)d_{q^2}s=\begin{cases}\frac{2}{1-q^2}\Theta_0 & for \quad z=1\\ 0 & for \quad z\neq 1.\end{cases}$$

Lemma 5.3

$$\mathcal{F}_{q^2}\Lambda=q^{-2}\Lambda^{-1}\mathcal{F}_{q^2},\qquad \mathcal{F}_{q^2}\partial_z=-is\mathcal{F}_{q^2},\qquad \mathcal{F}_{q^2}z=-iq^{-2}\Lambda^{-1}\partial_s\mathcal{F}_{q^2},\tag{5.4}$$

$$\mathcal{F}^{-1}_{q^2}\Lambda=q^{-2}\Lambda^{-1}\mathcal{F}^{-1}_{q^2},\qquad \mathcal{F}^{-1}_{q^2}\partial_s=i\Lambda^{-1}z\mathcal{F}^{-1}_{q^2},\qquad \mathcal{F}^{-1}_{q^2}s=i\partial_z\mathcal{F}^{-1}_{q^2}.\tag{5.5}$$

Proof. It is easy to verify that the kernels of the q^2-integral transforms (5.2) and (5.3) satisfy the conditions:

$${}_0\Phi_1(-;0;q^2,i(1-q^2)q^2zs)=\ddagger E_{q^2}(i(1-q^2)q^2zs)\ddagger,$$

$$E_{q^2}(-i(1-q^2)zs)=\ddagger e_{q^2}(i(1-q^2)zs)\ddagger$$

and

$$\partial_z\ddagger E_{q^2}((1-q^2)azs)\ddagger=a\ddagger E_{q^2}((1-q^2)aq^2zs)\ddagger s,$$

$$\partial_s\ddagger E_{q^2}((1-q^2)azs)\ddagger=az\ddagger E_{q^2}((1-q^2)aq^2zs)\ddagger,$$

$$\partial_z\ddagger e_{q^2}((1-q^2)azs)\ddagger=a\ddagger e_{q^2}((1-q^2)azs)\ddagger s,$$

$$\partial_s\ddagger e_{q^2}((1-q^2)azs)\ddagger=az\ddagger e_{q^2}((1-q^2)azs)\ddagger.$$

From (4.5) and the last relations we come to the statement of Lemma.■

Now we can prove Proposition 5.1.

From (5.4 we obtain

$$\mathcal{F}_{q^2} z^k \partial_z^l \phi(z) = (-i)^{k+l} q^{-2k} (\Lambda^{-1}\partial_s)^k s^l \mathcal{F}_{q^2}\phi(z).$$

On the other hand

$$\partial_s^k s^l = (-1)^k q^{k(2l-k+1)} \sum_{j=0}^{k} (-1)^j q^{j(j-1)} \frac{(q^{-2l}, q^2)}{(1-q^2)^{k-j}} \left[\begin{array}{c} k \\ j \end{array} \right]_{q^2} s^{l-k+j} \partial_s^j.$$

Therefore, if $\phi(z)$ satisfies (4.1) its image $\mathcal{F}_{q^2}\phi(z)$ satisfies (5.1). It means, in particular, that the image of converged sequence in $\hat{S}_{q^2}$ converges in $\hat{S}^{q^2}$. The statement concerning $\mathcal{F}_{q^2}^{-1}$ can be proved in the similar fashion using (5.5).

To prove the second part of Proposition we consider the action of the Fourier operators on the basis:

$$\mathcal{F}_{q^2} \circ \mathcal{F}_{q^2}^{-1} \hat{\psi}_n^{\pm}(s) = \hat{\psi}_n^{\pm}(s), \tag{5.6}$$

$$\mathcal{F}_{q^2}^{-1} \circ \mathcal{F}_{q^2} \hat{\phi}_n^{\pm}(z) = \hat{\phi}_n^{\pm}(z). \tag{5.7}$$

Consider the first relation

$$\mathcal{F}_{q^2} \circ \mathcal{F}_{q^2}^{-1} \hat{\psi}_n^{+}(s) = \frac{1}{2\Theta_0} \int d_{q^2} z \Big(\int E_{q^2}(-i(1-q^2)z\xi) \hat{\psi}_n^{+}(\xi) d_{q^2}\xi \Big) \, {}_0\Phi_1(-;0;q^2, i(1-q^2)q^2 zs) =$$

$$= \frac{1-q^2}{2\Theta_0} q^{2n} \int d_{q^2} z e_{q^2}(-i(1-q^2)q^{2n}z) \, {}_0\Phi_1(-;0;q^2, i(1-q^2)q^2 zs) =$$

$$= \frac{1-q^2}{2\Theta_0} \int d_{q^2} z e_{q^2}(-i(1-q^2)z) \, {}_0\Phi_1(-;0;q^2, i(1-q^2)q^{-2n+2} zs).$$

It follows from Lemma 5.1 that

$$\mathcal{F}_{q^2} \circ \mathcal{F}_{q^2}^{-1} \hat{\psi}_n^{+}(s) = \left\{ \begin{array}{lll} 1 & for & s = q^{2n} \\ 0 & for & s \neq q^{2n}. \end{array} \right.$$

and we come to (5.6). In the similar way Lemma 5.2 leads to (5.7).■

6 The q^2-Fourier transform of q^2-distributions

Definition 6.1 *The q^2-Fourier transform of q^2-distribution $f \in \hat{S}'_{q^2}$, is the q^2-distribution $g \in (\hat{S}^{q^2})'$ defined by the equality*

$$< g, \psi > = < f, \phi > \tag{6.1}$$

where $\phi(z)$ is an arbitrary function in S_{q^2} and $\psi(s) \in \hat{S}^{q^2}$ is its q^2-Fourier transform.

Suppose that the q^2-distribution f corresponds to $f(z)$ and $zf(z)$ is absolutely q^2-integrable function. Let $\phi(z) = \mathcal{F}_{q^2}^{-1}\hat{\psi}(s)$. Then

$$< f, \phi > = \frac{1}{2\Theta_0} \int d_{q^2} z \bar{f}(z) \int E_{q^2}(-i(1-q^2)zs) \psi(s) d_{q^2}s =$$

$$= \frac{1}{2\Theta_0}\int \overline{\int d_{q^2}z f(z)E_{q^2}(i(1-q^2)zs)}\psi(s)d_{q^2}s =< g,\psi > .$$

It means that the q^2-distribution g corresponds to the function

$$g(s) = \frac{1}{2\Theta_0}\int d_{q^2}z f(z)E_{q^2}(i(1-q^2)zs). \qquad (6.2)$$

Similarly, if g is defined by the absolutely q^2-integrable function $g(s)$ and $\hat{\psi}(s) = \mathcal{F}_{q^2}\hat{\phi}(z)$, then

$$< g,\psi >= \int\int d_{q^2}z\phi(z)\ {}_0\Phi_1(-;0;q^2,i(1-q^2)q^2zs)\bar{g}(s)d_{q^2}s =$$

$$= \int d_{q^2}z\phi(z)\overline{\int {}_0\Phi_1(-;0;q^2,-i(1-q^2)q^2zs)g(s)d_{q^2}s} =< f,\phi >,$$

i.e., f corresponds to

$$f(z) = \int {}_0\Phi_1(-;0;q^2,-i(1-q^2)q^2zs)g(s)d_{q^2}s. \qquad (6.3)$$

The q^2-Fourier transform of the q^2-distributions from $\hat{S}'_{q^2}$ we denote by $\mathcal{F}'_{q^2}$. The next Proposition follows from (6.2), (6.3) and Lemma 5.3

Proposition 6.1 *There are the following commutation relations in the space of q^2 distributions:*

$$\mathcal{F}'_{q^2}\Lambda = q^{-2}\Lambda^{-1}\mathcal{F}'_{q^2}, \qquad \mathcal{F}'_{q^2}\partial_z = -i\Lambda^{-1}s\mathcal{F}'_{q^2}, \qquad \mathcal{F}'_{q^2}z = -i\partial_s\mathcal{F}'_{q^2}. \qquad (6.4)$$

$$(\mathcal{F}'_{q^2})^{-1}\Lambda = q^{-2}\Lambda^{-1}(\mathcal{F}'_{q^2})^{-1}, (\mathcal{F}'_{q^2})^{-1}\partial_s = iz(\mathcal{F}'_{q^2})^{-1}, (\mathcal{F}'_{q^2})^{-1}s = iq^{-2}\Lambda^{-1}\partial_z(\mathcal{F}'_{q^2})^{-1}.$$

We define the q^2-Fourier transforms of some q^2-distributions.

1. It follows from (3.7) that

$$\mathcal{F}'_{q^2}z^{-1} = \frac{1}{2\Theta_0}\int d_{q^2}z z^{-1}E_{q^2}(i(1-q^2)zs) = \frac{i}{\Theta_0}\Theta(s).$$

Thus,

$$\mathcal{F}'_{q^2}z^{-1} = \hat{i}\,\mathrm{sign}\,s = i(\theta^+_{q^2} - \theta^-_{q^2}) \qquad (6.5)$$

as the distribution over $\hat{S}^{q^2}$.

2. (6.4), (6.5) and (3.7) give (4.9) that

$$\mathcal{F}'_{q^2}1 = \mathcal{F}'_{q^2}zz^{-1} = -i\partial_s\mathcal{F}'_{q^2}z^{-1} = \partial_s(\theta^+_{q^2} - \theta^-_{q^2}) = 2\delta_{q^2}. \qquad (6.6)$$

3.

$$\mathcal{F}'_{q^2}(\theta^+_{q^2} - \theta^-_{q^2}) = \frac{1}{2\Theta_0}\int_0^\infty d_{q^2}zE_{q^2}(i(1-q^2)zs) - \frac{1}{2\Theta_0}\int_{-\infty}^0 d_{q^2}zE_{q^2}(i(1-q^2)zs) =$$

$$= \frac{1-q^2}{2\Theta_0}\sum_{m=-\infty}^{\infty} q^{2m}[e_{q^2}(i(1-q^2)q^{2m}s) - e_{q^2}(-i(1-q^2)q^{2m}s)] =$$

$$= \frac{i(1-q^2)}{\Theta_0} \sum_{m=-\infty}^{\infty} q^{2m} \sin_{q^2}((1-q^2)q^{2m}s).$$

Using (2.15), we obtain

$$\mathcal{F}'_{q^2}(\theta^+_{q^2} - \theta^-_{q^2}) = \frac{is^{-1}}{\Theta_0} \lim_{M\to\infty} (1-q^2)s \sum_{m=-M}^{\infty} q^{2m} \sin_{q^2}((1-q^2)q^{2m}) =$$

$$= \frac{is^{-1}}{\Theta_0} \lim_{M\to\infty} [1 - \cos_{q^2}((1-q^2)q^{2M}] = \frac{is^{-1}}{\Theta_0}. \tag{6.7}$$

4. It follows from (4.9), (6.4) and (6.7) that

$$\mathcal{F}'_{q^2}\delta = \frac{1}{2}\mathcal{F}'_{q^2}\partial_z(\theta^+_{q^2} - \theta^-_{q^2}) = -\frac{i}{2}\Lambda^{-1}s\mathcal{F}'_{q^2}(\theta^+_{q^2} - \theta^-_{q^2}) = \frac{1}{2\Theta_0}.$$

5. From (4.7) and (4.8), (6.6) and (6.7) we obtain

$$\mathcal{F}'_{q^2}\theta^+_{q^2} = \frac{1}{2}\mathcal{F}'_{q^2}(\theta^+_{q^2} - \theta^-_{q^2} + 1) = \frac{is^{-1}}{2\Theta_0} + \delta_{q^2},$$

$$\mathcal{F}'_{q^2}\theta^-_{q^2} = \frac{1}{2}\mathcal{F}'_{q^2}(-\theta^+_{q^2} + \theta^-_{q^2} + 1) = -\frac{is^{-1}}{2\Theta_0} + \delta_{q^2}.$$

Proposition 6.2 *For an arbitrary integer* $n \geq 0$

$$\mathcal{F}'_{q^2} z^n = 2i^n q^{-n(n+1)} \frac{(q^2,q^2)_n}{(1-q^2)^n} s^{-n} \delta_{q^2}(s).$$

Proof. The relation $1 = \frac{(1-q^2)^n}{(q^2,q^2)_n}\partial_z^n z^n$ follows from (3.2). Next, from (6.6) and Proposition 6.1 we obtain

$$\delta_{q^2}(s) = \frac{1}{2}\mathcal{F}'_{q^2}1 = \frac{(1-q^2)^n}{2(q^2,q^2)_n}\mathcal{F}'_{q^2}\partial_z^n z^n = (-i)^n \frac{(1-q^2)^n}{2(q^2,q^2)_n}(\Lambda^{-1}s)^n \mathcal{F}'_{q^2} z^n.$$

Thus,

$$\mathcal{F}'_{q^2} z^n = 2i^n \frac{(q^2,q^2)_n}{(1-q^2)^n} (s^{-1}\Lambda)^n \delta_{q^2}(s). \tag{6.8}$$

Since $(s^{-1}\Lambda)^n = q^{-n(n-1)} s^{-n} \Lambda^n$ and $\delta_{q^2}(q^{2n}s) = q^{-2n}\delta_{q^2}(s)$,

$$(s^{-1}\Lambda)^n \delta_{q^2}(s) = q^{-n(n+1)} s^{-n} \delta_{q^2}(s).$$

Then, the statement of Proposition follows from (6.8). ■

Proposition 6.3 *For an arbitrary integer* $n \geq 0$

$$\mathcal{F}'_{q^2} z^{-n-1} = i^{n+1} \frac{(q^2,q^2)_n}{(1-q^2)^n} s^n \operatorname{sign} s.$$

Proof. For an arbitrary $n \geq 0$ we have from (3.3)

$$\partial_z^n z^{-1} = (-1)^n q^{-n(n+1)} \frac{(q^2, q^2)_n}{(1-q^2)^n} z^{-n-1}.$$

Using Proposition 6.1 we obtain

$$\mathcal{F}'_{q^2} z^{-n-1} = (-1)^n q^{n(n+1)} \frac{(1-q^2)^n}{(1-q^2)^n} \mathcal{F}'_{q^2} \partial_z^n z^{-1} = i^n q^{n(n+1)} \frac{(1-q^2)^n}{(1-q^2)^n} (\Lambda^{-1} s)^n \mathcal{F}'_{q^2} z^{-1}.$$

It can be derived from (4.5) that $(\Lambda^{-1}s)^n = q^{-n(n+1)} s^n \Lambda^{-n}$. Using this equality and (6.5), we come to the statement of Proposition. ∎

Proposition 6.4

$$\mathcal{F}'_{q^2} z_+^{\nu-1} = \frac{e_{q^2}(q^2) E_{q^2}(-q^{2(1-\nu)})}{2\Theta_0} (\bar{c}_\nu s_-^{-\nu} + c_\nu s_+^{-\nu}), \tag{6.9}$$

$$\mathcal{F}'_{q^2} z_-^{\nu-1} = -\frac{e_{q^2}(q^2) E_{q^2}(-q^{2(1-\nu)})}{2\Theta_0} (c_\nu s_-^{-\nu} + \bar{c}_\nu s_+^{-\nu}), \tag{6.10}$$

where

$$c_\nu = \sum_{m=-\infty}^{\infty} \frac{q^{2\nu m}(q^{-2m} + i(1-q^2))}{(1-q^2)^{-1} q^{-2m} + (1-q^2) q^{2m}}.$$

Proof. If $0 < \mathrm{Re}\, \nu < 1$, (4.10) and (6.2) give

$$\mathcal{F}_{q^2} z_+^{\nu-1} = \frac{1-q^2}{2\Theta_0} \sum_{m=-\infty}^{\infty} q^{2\nu m} e_{q^2}(i(1-q^2) q^{2m} s). \tag{6.11}$$

It can be directly checked that $z_+^{\nu-1}$ satisfies the equation

$$z \partial_z f(z) = \frac{1-q^{2(\nu-1)}}{1-q^2} f(z).$$

From the Proposition 6.1 we obtain that the q^2-Fourier transformed equation takes the form

$$-q^{-2} \partial_s (s g(q^{-2} s)) = \frac{1-q^{2(\nu-1)}}{1-q^2} g(s).$$

Since $s_-^{-\nu}$ and $s_+^{-\nu}$ satisfy this equation we can write

$$\mathcal{F}'_{q^2} z_+^{\nu-1} = c_1 s_-^{-\nu} + c_2 s_+^{-\nu}. \tag{6.12}$$

Substituting $s = 1$ in (6.12) we obtain from (2.6),(6.11) and (6.12)

$$c_2 = \frac{1-q^2}{2\Theta_0} \sum_{m=-\infty}^{\infty} q^{2\nu m} e_{q^2}(i(1-q^2) q^{2m}) =$$

$$= \frac{1-q^2}{2\Theta_0} e_{q^2}(q^2) \sum_{m=-\infty}^{\infty} q^{2\nu m} \sum_{n=0}^{\infty} \frac{(-1)^n q^{n(n+1)}}{(q^2, q^2)_n [1 - i(1-q^2) q^{2(m+n)}]}.$$

If $0 < \mathrm{Re}\,\nu < 1$, we can change the order of summations. Then

$$c_2 = \frac{1-q^2}{2\Theta_0} e_{q^2}(q^2) \sum_{n=0}^{\infty} \frac{(-1)^n q^{n(n+1)}}{(q^2,q^2)_n} \sum_{m=-\infty}^{\infty} \frac{q^{2\nu m}}{1 - i(1-q^2)q^{2(m+n)}} =$$

$$= \frac{e_{q^2}(q^2)}{2\Theta_0} \sum_{n=0}^{\infty} \frac{(-1)^n q^{n(n-1)} q^{2(1-\nu)n}}{(q^2,q^2)_n} \sum_{m=-\infty}^{\infty} \frac{q^{2\nu m}(q^{-2m} + i(1-q^2))}{(1-q^2)^{-1} q^{-2m} + (1-q^2) q^{2m}}.$$

We denote the sum of the inner series by c_ν. Then

$$c_2 = \frac{c_\nu}{2\Theta_0} e_{q^2}(q^2) E_{q^2}(-q^{2(1-\nu)})$$

Assume now that $s = -1$ in (6.12). Then in the same way we obtain

$$c_1 = \frac{\bar{c}_\nu}{2\Theta_0} e_{q^2}(q^2) E_{q^2}(-q^{2(1-\nu)})$$

and thereby come to (6.9). It determines $\mathcal{F}'_{q^2} z_+^{\nu-1}$ as (6.9) for non-integer ν by the analytical continuation.

The q^2-Fourier transform of $z_-^{\nu-1}, \quad \nu \neq 0, \pm 1, \ldots$.

$$\mathcal{F}'_{q^2} z_-^{\nu-1} = -\frac{1-q^2}{2\Theta_0} \sum_{m=-\infty}^{\infty} q^{2\nu m} e^{q^2}(-i(1-q^2)q^{2m}s).$$

differs from (6.11) by the sign and the complex conjugation, i.e., in this way we have (6.10). ■

We summarize all results in the table:

$f(z)$	$g(s) = \frac{1}{2\Theta_0} \int d_{q^2} z f(z) E_{q^2}(i(1-q^2)zs)$
$\delta_{q^2}(z)$	$\frac{1}{2\Theta_0}$
$\theta^+_{q^2}(z)$	$\frac{i}{2\Theta_0} s^{-1} + \delta_{q^2}(s)$
$\theta^-_{q^2}(z)$	$-\frac{i}{2\Theta_0} s^{-1} + \delta_{q^2}(s)$
z^n $(n = 0, 1, \ldots)$	$2 i^n q^{-n(n+1)} \frac{(q^2,q^2)_n}{(1-q^2)^n} s^{-n} \delta_{q^2}(s)$
z^{-n-1} $(n = 0, 1, \ldots)$	$i^{n+1} \frac{(1-q^2)^n}{(q^2,q^2)_n} s^n \,\mathrm{sign}\, s$
$z_+^{\nu-1}$ $(\nu \neq 0, \pm 1, \ldots)$	$\frac{e_{q^2}(q^2) E q^2(-q^{2(1-\nu)})}{2\Theta_0} (\bar{c}_\nu s_-^{-\nu} + c_\nu s_+^{-\nu})$
$(z_-^{\nu-1}$ $(\nu \neq 0, \pm 1, \ldots)$	$-\frac{e_{q^2}(q^2) E q^2(-q^{2(1-\nu)})}{2\Theta_0} (c_\nu s_-^{-\nu} + \bar{c}_\nu s_+^{-\nu}]$

Acknowledgments.

The work is supported in part by grants 96-15-96455 for support of scientific schools, INTAS-93-0166 extension and by Award No. RM2-150 of the Civilian Research & Development Foundation (CRDF) for the Independent States of the Former Soviet Union (M.O.); by RFBR 97-01-00747 and NIOKR MPS RF (V.R.) The work is prepared partly during visit of M.O. in IHES (Bur-sur-Yvette). He is grateful to Prof. K.Gawedzky for hospitality.

References

[1] Chryssomalakos Ch. and Zumino B.,*Translations, Integrals and Fourier Transforms in the Quantum Plane.* Salamfest, 1993, 327-346

[2] Kempf A. and Majid S., *Algebraic q-Integration and Fourier Theory on Quantum and Braided Spaces*, J.Math.Phys. 35 (1994) 6802-6837

[3] Koornwinder T. and Swarttouw R., q-analogues of the Fourier and Hankel Transforms. Trans. AMS. **333**, 1 (1992) 445-461

[4] Askey, R., Atakisheiev, N. and Suslov, S *An analog of the Fourier transformation for the q-harmonic oscillator*, Symmetries in Science VI (1993) 57-63 (B.Gruber, ed.) Plenum Press, NY

[5] Rahman, M. and Suslov, S. *Singular analogue of the Fourier transformation for the Askey-Wilson polynomials*, Preprint CRM-1915, (1993)

[6] Askey, R., Rahman, M., and Suslov, S., *On a general q-Fourier transformation with non-symmetric kernels*, Preprint ISSN 0827-3669 (1994)

[7] Gel'fand I.M. and Shilov G.E.,*Generalized functions,* Vol.1: Properties and operations., M., Fizmatgiz, 1959; English transl. of 1st ed., Academic Press, 1964

[8] T. H. Koornwinder, *Special functions and q-commuting variables*, in "Special Functions, q-Series and Related Topics", M. E. H. Ismail, D. R. Masson and M. Rahman (eds.), Fields Institute Communications 14, American Mathematical Society, 1997, pp. 131–166.

[9] Gasper G. and Rahman M.,*Basic Hypergeometric Series,* Cambridge: Cambridge University Press, 1990